

The Examination Paper of Jinan University (Solutions)

For Instructor Only	Academic Year: <u>2016–2017</u>	Semester: 1 st [<input type="checkbox"/>] 2 nd [<input checked="" type="checkbox"/>]	Course Type Compulsory [<input checked="" type="checkbox"/>] Elective [<input type="checkbox"/>]
	Course Title: _____	Math 1906	
	Instructor's Name: _____	Jianrui Lyu	Form of the Examination Open-book [<input type="checkbox"/>] Closed-book [<input checked="" type="checkbox"/>]
	Date of the Examination: _____	dd/mm/yyyy	Paper A [<input checked="" type="checkbox"/>] Paper B [<input type="checkbox"/>] Total Pages <u>6</u>
For Student Only	School/College _____	Major _____	
	Name _____	Student No. _____	
Mainland Student [<input type="checkbox"/>]	Non-mainland Student [<input checked="" type="checkbox"/>]		

Section No.	I	II	III	IV	Total Score
Score					

Section I: Fill in the blanks. (6 questions; 3 points for each; 18 points in total.)

Notice: you MUST write the answers in the following tables.

Number	1	2	3
Answer			
Number	4	5	6
Answer			

Score	Evaluator	Section II: Select one answer from four choices. (6 questions; 3 points for each; 18 points in total.)

Notice: you MUST write the answers in the following tables.

Number	1	2	3	4	5	6
Answer						

1. The first question text, text (C)
 (A) first $\int f'(x)dx = f(x)$ (B) second $\int df(x) = f(x)$
 (C) third $\frac{d}{dx}(\int f(x)dx) = f(x)$ (D) fourth $d(\int f(x)dx) = f(x)$

2. The second question $F(x)$ text $f(x)$ text, text (A)
 (A) first choice $F(x)$ text $\Leftrightarrow f(x)$ text
 (B) second choice $F(x)$ text text $\Leftrightarrow f(x)$ text
 (C) third choice $F(x)$ text $\Leftrightarrow f(x)$ text text
 (D) fourth choice $F(x)$ text text $\Leftrightarrow f(x)$ text text

3. The third question $A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & x & 0 \\ 0 & 0 & 1 \end{pmatrix}$ text text $\lambda_1 = 1$ text $\lambda_2 = 2$, text $x =$ (B)
 (A) 2 (B) 1 (C) 0 (D) -1

4. The fourth question $f = 4x_1^2 - 2x_1x_2 + 6x_2^2$ text (C)
 (A) $\begin{pmatrix} 4 & -2 \\ -2 & 6 \end{pmatrix}$ (B) $\begin{pmatrix} 2 & -2 \\ -2 & 3 \end{pmatrix}$ (C) $\begin{pmatrix} 4 & -1 \\ -1 & 6 \end{pmatrix}$ (D) $\begin{pmatrix} 2 & -1 \\ -1 & 3 \end{pmatrix}$

5. The fifth question wrong text (B)
 (A) first choice text text text text text text text text text
 (B) second choice text text text text text text text text text
 (C) third choice text text text text text text text text text
 (D) fourth choice text text text text text text text text

6. The sixth question X text (X_1, \dots, X_n) text wrong text (D)
 (A) text text text (B) text n text (C) X_1, \dots, X_n text (D) $X_1 = X_2 = \dots = X_n$

Score	Evaluator	Section III: Work out math questions. (6 questions; 8 points for each; 48 points in total.)

1. The first question $\int e^{2x} (\tan x + 1)^2 dx$.

Solution. $I = \int e^{2x} \sec^2 x dx + 2 \int e^{2x} \tan x dx$ 2 points
 $= \int e^{2x} d(\tan x) + 2 \int e^{2x} \tan x dx$ 4 points
 $= e^{2x} \tan x - 2 \int e^{2x} \tan x dx + 2 \int e^{2x} \tan x dx$ 6 points
 $= e^{2x} \tan x + C$ 8 points

2. The second question $A(1, 2, -1), B(2, 3, 0), C(3, 3, 2)$ text $\triangle ABC$ text text text text text text.

Solution. Text $\overrightarrow{AB} = (1, 1, 1), \overrightarrow{AC} = (2, 1, 3)$, 2 points
 text $\overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 1 & 1 \\ 2 & 1 & 3 \end{vmatrix} = (2, -1, -1)$, 4 points
 text $\triangle ABC$ text $S_{\triangle ABC} = \frac{1}{2} |\overrightarrow{AB} \times \overrightarrow{AC}| = \frac{1}{2} \sqrt{6}$ 6 points
 Text text $2(x-2)-(y-3)-z=0$, text $2x-y-z-1=0$ 8 points

3. The third question $A = \begin{vmatrix} 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 0 \\ 2 & 3 & 0 & 1 \\ 3 & 0 & 1 & 2 \end{vmatrix}$ text.

$$\begin{aligned}
 \text{Solution. } A &= \left| \begin{array}{cccc} 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 0 \\ 2 & 3 & 0 & 1 \\ 3 & 0 & 1 & 2 \end{array} \right| = \left| \begin{array}{cccc} 0 & 1 & 2 & 3 \\ 1 & 2 & 3 & 0 \\ 0 & -1 & -6 & 1 \\ 0 & -6 & -8 & 2 \end{array} \right| = 1 \cdot (-1)^{2+1} \left| \begin{array}{ccc} 1 & 2 & 3 \\ -1 & -6 & 1 \\ -6 & -8 & 2 \end{array} \right| \quad \dots\dots\text{4 points} \\
 &= - \left| \begin{array}{ccc} 1 & 2 & 3 \\ 0 & -4 & 4 \\ 0 & 4 & 20 \end{array} \right| = - \left| \begin{array}{cc} -4 & 4 \\ 4 & 20 \end{array} \right| = -(-4 \cdot 20 - 4 \cdot 4) = 96 \quad \dots\dots\text{8 points}
 \end{aligned}$$

4. The fourth question, $f = x_1^2 + 2x_1x_2 - 6x_1x_3 + 2x_2^2 - 12x_2x_3 + 9x_3^2$ $f = d_1y_1^2 + d_2y_2^2 + d_3y_3^2$.

Solution. $f = x_1^2 + 2x_1x_2 - 6x_1x_3 + 2x_2^2 - 12x_2x_3 + 9x_3^2$

$$= x_1^2 + 2x_1(x_2 - 3x_3) + (x_2 - 3x_3)^2 + x_2^2 - 6x_2x_3 \quad \dots \text{3 points}$$

$$= (x_1 + x_2 - 3x_3)^2 + x_2^2 - 6x_2x_3$$

$$= (x_1 + x_2 - 3x_3)^2 + x_2^2 - 2x_2 \cdot 3x_3 + (3x_3)^2 - 9x_3^2 \quad \dots \text{6 points}$$

$$= (x_1 + x_2 - 3x_3)^2 + (x_2 - 3x_3)^2 - 9x_3^2$$

Text $y_1 = x_1 + x_2 - 3x_3$, $y_2 = x_2 - 3x_3$, $y_3 = x_3$,

text $f = y_1^2 + y_2^2 - 9y_3^2$ text. \dots 8 points

- 5.** The fifth question text text text 0.2 text text, text text 100 text text.

(1) text text text text text text ξ text 10 text 30 text.

(2) text text text text text ξ text 10 text 30 text.

Solution. $E\xi = np = 100 \cdot 0.2 = 20$, $D\xi = npq = 100 \cdot 0.2 \cdot 0.8 = 16$ 2 points

$$(1) P(10 < \xi < 30) = P(|\xi - E\xi| < 10) \geq 1 - \frac{D\xi}{10^2} = 1 - \frac{16}{100} = 0.84. \quad \dots\dots \text{4 points}$$

$$(2) P(10 < \xi < 30) \approx \Phi_0\left(\frac{30-20}{\sqrt{16}}\right) - \Phi_0\left(\frac{10-20}{\sqrt{16}}\right) \quad \dots \text{.....} \textcolor{red}{6 \text{ points}}$$

$$= 2\Phi_0(2.5) - 1 = 2 \cdot 0.9938 - 1 = 0.9876 \quad \dots\dots \text{8 points}$$

6. The sixth question $N(\mu, \sigma^2)$ text text 16 text, text text text 3160, text text 100. Text text $H_0 : \mu = 3140$ text text ($\alpha = 0.01$).

Solution. (1) Text text $H_0 : \mu = 3140$ 2 points

(2) Text text text $T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$ 3 points

(3) Text text $t_a = t_a(n-1) = t_{0.01}(15) = 2.947$ 5 points

$$(4) \text{Text text text } t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}} = \frac{3160 - 3140}{100/4} = 0.8. \quad \dots\dots \text{7 points}$$

(5) Text $|t| < t_a$, text text H_0 , text text text. 8 points

(3) $\lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n = e$, $\lim_{n \rightarrow \infty} n! = \infty$.

Score	Evaluator	Section IV: Work out math proofs. (2 questions; 16 points in total.)

1. (9 points) The first question $\{x_n\}$ text $x_1 = \sqrt{2}$, $x_{n+1} = \sqrt{2 + x_n}$. Text text text, text text text.

Proof. (1) Text, text $x_1 < 2$, text $x_k < 2$ text

$$x_{k+1} = \sqrt{2 + x_k} < \sqrt{2 + 2} = 2,$$

Text text text text text n text $x_n < 2$, text text text. Text text

$$\frac{x_{n+1}}{x_n} = \sqrt{\frac{2}{x_n^2} + \frac{1}{x_n}} > \sqrt{\frac{2}{2^2} + \frac{1}{2}} = 1,$$

Text text text text text. Text text text text text, Text text text text.

..... 4 points

(2) Text text text text *A*, text text text text text text

$$A = \sqrt{2 + A}.$$

Text text $A = 2$, text text $\{x_n\}$ text text text 2.

..... 8 points

2. (7 points) The second question A text B text, text A text \bar{B} text.

Proof. $P(A \cdot \bar{B}) = P(A - B) = P(A - AB)$

..... 2 points

$$= P(A) - P(AB) = P(A) - P(A)P(B)$$

..... 4 points

$$= P(A)(1 - P(B)) = P(A)P(\bar{B})$$

.....6 points

Text text text A text text text \bar{B} text text text.

.....8 points

Appendix Some data may be used in the exam

$$\begin{array}{cccc} \Phi_0(0.5) = 0.6915 & \Phi_0(1) = 0.8413 & \Phi_0(2) = 0.9773 & \Phi_0(2.5) = 0.9938 \\ t_{0.01}(8) = 3.355 & t_{0.01}(9) = 3.250 & t_{0.01}(15) = 2.947 & t_{0.01}(16) = 2.921 \\ \chi^2_{0.005}(8) = 22.0 & \chi^2_{0.005}(9) = 23.6 & \chi^2_{0.005}(15) = 32.8 & \chi^2_{0.005}(16) = 34.3 \\ \chi^2_{0.995}(8) = 1.34 & \chi^2_{0.995}(9) = 1.73 & \chi^2_{0.995}(15) = 4.60 & \chi^2_{0.995}(16) = 5.14 \end{array}$$